

Taq DNA Polymerase

Cat. No. TAQ001

Store at -20°C.

Product Description

Taq DNA Polymerase is a highly thermostable DNA Polymerase that catalyzes the 5'-3' synthesis of DNA. This polymerase has 5'-3' exonuclease activities, lacks 3'-5' exonuclease activity, and produces 3'-dA-tailed amplicons. PCR products made with Taq can be used with TA cloning vectors.

Product Component	Quantity	Part No.
Taq DNA Polymerase	200 rxn (200 µl)	TAQ001.200
5X Taq Buffer ¹	2 x 1.0 ml	TAQ001.B

¹. Buffer contains 1.5 mM Mg²⁺

Protocol

- Mix individual components before use and assemble reaction on ice.

Component	Volume
5X Taq Buffer	10 µl
dNTP Mix (10 mM)	1 µl
Forward Primer (10 µM)	1-2 µl
Reverse Primer (10 µM)	1-2 µl
Template DNA	Variable (200 ng genomic DNA)
Taq DNA Polymerase	1 µl
Nuclease-free H ₂ O	up to 50 µl

- Gently mix the reaction components and briefly centrifuge. Run thermocycling conditions for standard PCR:

Step	Temperature	Duration
Initial Denaturation ²	94°C	3 min
25 – 35 Cycles	94° C	30 sec
	50-65° C ³	30 sec
	72° C	1 min/kb
Final Extension	72° C	5 min

²For most applications, an initial 3 minute denaturation step at 94°C is sufficient. Increase to 5 minutes for high-GC or difficult templates.

³Adjust annealing temperature based on the primer melting temperature.

General Notes

- For difficult or high-GC templates, use a standard touchdown program.
- If working with low template and/or low sample amount, reduce total reaction volume to 25 µl and lower other component volumes accordingly.

BioShop Canada Inc.

5480 Mainway, Burlington, Ontario, Canada, L7L 6A4

Phone: 1-800-234-1179 Fax: 1-905-634-7356 Email: sales@bioshopcanada.com